

Index

| - A - | |
|---|---|
| absorption | A3.3 |
| absorption coefficient | F4.3 |
| absorption correction | D3.9 |
| absorption edge | D11.9 |
| absorption length | A3.2 |
| acoustic | B4.5, E7.8, E7.11 |
| adiabatic approximation | B4.2 |
| adiabatic potential energy surface | B5.19 |
| a-Helix | E10.6 |
| alcohol dehydrogenase | E3.19, E7.9 |
| alpha- and beta-relaxation | D9.15, D9.16 |
| amino acid | E10.2 |
| amorphous | B4.10 |
| amphiphilic block copolymers | B2.25 |
| amphiphilic systems | B2.2 |
| analyzer crystal | C7.17, C7.19, C7.20, C7.26, C7.27, C7.30 |
| anchored polymers | B2.26 |
| anharmonic interaction potential | E4.9 |
| anharmonicity | B4.8, E7.9, E9.3, E9.5, E9.13, E9.14, E9.18 |
| anisotropy of tensor of susceptibility | D11.7 |
| anomalous dispersion | A4.11, E10.15 |
| anomalous scattering | A1.18, D11.3 |
| anomalous small angle x-ray scattering | D1.22 |
| anomalous transmission | A3.3 |
| antiferromagnet | B4.14, C7.8, C7.34, E5.9 |
| atomic disorder | D3.25 |
| atomic form factor | A4.7 |
| atomic Pair Distribution Function (PDF) | D5.2, D5.3, E7.11 |
| atomic vibrations | E7.8, E7.9, E7.10, E7.11, E7.16, E8.9 |
| ATS | D11.7 |
| attenuation coefficient | F2.5 |
| autocorrelation function | D10.9 |
| azimuth dependence | D11.8 |
| - B - | |
| backscattering spectrometer | C3.22, D7.2, E7.13 |
| balanced microemulsions | B2.24 |
| band structure | A3.8 |
| bead modeling | E8.5 |
| beamtime proposal | A1.39 |
| bending modulus | D8.9 |
| bending rigidity | B2.9 |
| bicontinuous microemulsion | B2.6 |
| bicontinuous structure | B2.6 |
| BiFeO ₃ | E4.12 |
| binary amphiphilic systems | B2.3 |

| bio-diffractometer | C3.10 |
|---|-------------------------|
| biological macromolecules | E8.2 |
| biology | B6.2 |
| biomimetic membranes | E5.7 |
| Bloch's theorem | A3.3, B4.4 |
| block copolymer | E5.3 |
| Blume-Maleyev equations | C6.15, C6.19 |
| Bogoliubov transformation | B4.16 |
| Born approximation | A1.7, A3.2, D1.5, D1.10 |
| Born series | A3.12 |
| Born-Oppenheimer approximation | B5.18 |
| Borrmann effect | A3.3 |
| Bose factor | D4.14 |
| Bragg condition | A5.8 |
| Bragg diffraction | A3.3 |
| Bragg equation | A1.13 |
| Bragg scattering | A3.4 |
| Bragg's law | D3.4 |
| Brockhouse | A1.3 |
| b-sheet | E10.6 |
| buffer | E8.2 |
| - C - | |
| Ca ₃ CoMnO ₆ | E4.16 |
| Chadwick | A1.3 |
| charge order | D11.12, E4.16 |
| chemical shifts | F3.20 |
| circular dichroism | F4.19 |
| classical approximation | A5.19 |
| cloning theory | E9.15, E9.18 |
| Co ₈₀ Fe ₂₀ | E5.9 |
| coherence | A1.9, D1.9 |
| coherence length (transverse, longitudinal) | D8.15 |
| coherence volume | D2.8 |
| coherence, longitudinal | C8.13 |
| coherence, transverse | C8.3, C8.7 |
| coherent diffraction | D10.4 |
| coherent diffraction imaging | C8.7 |
| coherent neutron scattering cross section | A4.14 |
| coherent scattering | A5.15 |
| cold neutrons | C1.12 |
| collimation ratio | F2.4 |
| collimator | C3.3 |
| colloidal suspensions | B2.2 |
| colloids (charge stabilized) | D8.14 |
| colors of transition-metal complexes | B5.8 |
| complex fluids | B2.2, D1.2 |
| Compton scattering | A4.6, D5.5, D5.7, D5.10 |
| continuous slowing down approximation | A4.12 |

| | |
|---------------------------------------|---|
| contour length fluctuations | E3.16 |
| contrast | D2.13, D2.17, D2.19, D2.25 |
| contrast enhancement | E5.7 |
| contrast variation | D1.18, E6.2, E6.7, E6.8, E6.10, E6.12, E6.13, E6.15, E6.16, E7.14 |
| convolution | A5.29 |
| cooperatively rearranging regions | E9.11, E9.15 |
| coordination compounds | B5.2 |
| correlation function | A5.30, D6.7, D6.13-D6.17, E7.13 |
| correlation length | B2.15, D1.17, D6.4, D6.5, D6.7, D6.10, D6.17-D6.21, D8.2 |
| coupling model | E9.16, E9.18, E9.20 |
| critical dynamics | D6.13 |
| critical exponents | D6.6-D6.9, D6.11-D6.13, D6.21 |
| critical scattering | D6.3, D6.17-D6.21 |
| critical temperature | D6.4 |
| cross section | A1.7 |
| cryptotomography | D10.18 |
| crystal field | B5.3 |
| crystal field splitting | B5.6 |
| crystal lattice | B1.2 |
| crystal structure | B1.2 |
| crystalline lattice | A5.7 |
| crystallography | D3.3 |
| cultural heritage studies | F2.20 |
| curvature elasticity | B2.9 |
| curvature energy | B2.9 |
| cytoskeleton | B6.2 |
| - D - | |
| dammin | E8.6 |
| Darwin profile | A3.18 |
| Darwin width | A3.18 |
| Debye model | D7.4 |
| Debye-Waller factor | A5.26, D4.13, D7.4 |
| delta function | A5.28 |
| density matrix | C7.11, C7.12, C7.26 |
| detector | C3.7 |
| differential cross-section | A5.3 |
| differential PDF | D5.8 |
| differential scattering cross section | A4.2 |
| diffraction | A3.2 |
| diffraction before destruction | D10.3 |
| diffuse scattering | D5.2, D5.3, D5.9 |
| diffusion | B3.1, D7.6, D9.2, D9.11, D9.12, D9.13, E7.12 |
| diffusion equations | B3.7, B3.9, B3.17 |
| dipole approximation | F3.5 |
| dipole selection rules | F3.40 |
| discovery neutron | F2.2 |
| disordered systems | D9.15 |
| dispersion equation | A3.8 |

| | |
|---|--|
| dispersion surface | A3.7 |
| displacive phase transition | B4.9 |
| dissipation-fluctuation theorem | D4.12 |
| disulfide bridge | E10.5 |
| diverging length scale | E9.15 |
| DKD | C3.13 |
| DNA | B6.2 |
| DOS | D9.8, D9.9, D9.11, D9.12, D9.15, D9.16 |
| double differential cross section | C3.2, D4.12 |
| double differential cross-section | A5.12 |
| DVD | E4.4 |
| Dynamic Light Scattering (DLS) | D8.11 |
| dynamic magnetic susceptibility | D6.19 |
| dynamic modulus | E3.4 |
| dynamic scattering | A5.11 |
| dynamic structure factor | B6.16 |
| dynamical beats | D9.3, D9.4, D9.5 |
| dynamical heterogeneity | E9.1, E9.7, E9.8, E9.9, E9.16 |
| dynamical matrix | B4.4 |
| dynamical scattering | A3.2 |
| dynamical structure factor | A1.26, D4.13 |
| dynamically triangulated surfaces | B2.20 |
| DWBA | D2.30 |
| Dzyaloshinskii-Moriya interaction | E4.14 |
| - E - | |
| efficiency boosting, amphiphilic block copolymers | B2.25 |
| eigenfrequency | B4.4 |
| eigenfunction | B6.7 |
| eigenfunction expansion | B6.6 |
| eigenvalue | B6.6 |
| eigenvector | B4.4 |
| Einstein | F3.2 |
| elastic incoherent structure factor | D7.4 |
| elastic scattering | A5.3 |
| electron scattering | A4.12 |
| electron spin resonance | B5.15 |
| electronic correlations | F3.30 |
| electron-magnon interaction | F3.33 |
| electron-phonon interaction | F3.32 |
| Eliashberg function | F3.32 |
| emittance | C8.2 |
| emulsification failure | B2.12 |
| energy distribution curve | F3.18 |
| energy landscape | E9.17, E9.18, E9.20 |
| energy selection | F2.16 |
| ensemble average | D8.11 |
| entanglements | B6.19, E3.10 |

| | |
|--|--------------------------------|
| entropy | B6.2 |
| envelope function | D5.9, D5.11 |
| Error Reduction algorithm | D10.11 |
| ESCA | F3.3 |
| eukaryotic cells | B6.2 |
| Ewald construction | D3.5 |
| Ewald sphere | A3.7, E10.14 |
| EXAFS (Extended X-ray Absorption Fine Structure) | F4.4, F4.17 |
| exchange bias | E4.10, E5.9 |
| exchange splitting | F3.25 |
| exchange-correlation potential | F3.4 |
| excluded volume | B6.15 |
| extinction correction | D3.10 |
| extinction length | A3.2 |
| - F - | |
| Fano effect | F4.22 |
| Fano resonances | F4.11 |
| FEL (Free-electron Laser) | C8.1 |
| Fermi | A1.3 |
| Fermi pseudopotential | A4.14 |
| Fermi surface mapping | F3.21, F3.24 |
| Fermi's Golden Rule | F3.4, F4.5 |
| ferroelectrics | D6.3, D6.5, D6.21 |
| ferroic order | E4.8 |
| ferromagnet | B4.12, C7.34, C7.37, E5.9 |
| Fick's diffusion equation | B3.7 |
| first cumulant expansion | E3.19 |
| fission | A1.16 |
| FLASH | C8.5 |
| flat-cone | C3.8 |
| flexible polymer | B6.5 |
| flipping ratio | C7.34, C7.36 |
| Flory exponent | D1.16 |
| fluorescence correlation spectroscopy (FCS) | B6.14 |
| focusing instrument | D1.7 |
| forbidden reflections | D11.5 |
| form factor | A1.18, A5.10, D3.7 |
| Fourier time | C3.27 |
| Fourier transformation | A5.26, D1.5 |
| fragility | E9.1, E9.6, E9.7, E9.11, E9.16 |
| free electron laser | A1.15 |
| free-draining | B6.6 |
| free-electron laser, FEL | D10.2 |
| frustration | E4.14 |
| full width at half maximum | D7.6 |
| - G - | |
| gain length | C8.7 |
| Gaussian approximation | E3.12 |
| Gaussian fluctuations | B2.31 |
| Gaussian free-energy functional | B2.31 |
| Gaussian polymer | B6.3 |
| Gaussian process | B6.6 |
| Gaussian random fields | B2.15, B2.16, B2.32 |

| | |
|---|---|
| Ge2Sb2Te5 | E4.3 |
| geometrical averages | B2.16, B2.32 |
| GeTe | E4.5 |
| giant magnetoresistance effect | E4.9 |
| Gibbs-Bogoliubov-Feynman inequality | B2.17 |
| Ginzburg-Landau model | D6.4, D6.9, D6.10, D6.12 |
| Ginzburg-Landau model, fluctuations | D2.14, B2.31 |
| Ginzburg-Landau theory | D6.4, D6.5, D6.11 |
| GISAXS, GISANS | A1.34, B2.29, C3.15, D2.3, D2.6, D2.7, D2.36, E5.3, E7.14 |
| glass | D5.2, D5.5, D5.10, D9.15, D9.16, D9.17, E7.9 |
| glass transition | E9.1 ff |
| gradient diffusion | B3.2, B3.15 |
| Green's function | A3.12 |
| Guinier scattering | D1.14 |
| - H - | |
| harmonic approximation | B4.2 |
| harmonic oscillator | A5.24, B6.4 |
| Heisenberg model | B4.12, D6.8, D6.9 |
| Helfrich-model | D8.10 |
| heterostructure | E5.1 |
| hexaferrite | E4.16 |
| high pressure | D9.2, D9.8, D9.9, D9.10 |
| high-spin complex | B5.10 |
| Holstein-Primakoff transformation | B4.13 |
| hot neutrons | C1.11 |
| Hybrid Input-Output algorithm (HIO) | D10.11 |
| hydrodynamic interactions | B6.9 |
| hydrogen | D7.2, E7.12, E7.13 |
| hydrogen bond | E10.5 |
| hydrogen diffusion in metals | B3.20 |
| hydrophobic cluster | E10.5 |
| hydrophobic-hydrophilic-balance temperature | B2.6 |
| hyperfine spectroscopy | D9.2 |
| Hyperscaling | D6.11 |
| hysteresis | E5.10 |
| - I - | |
| ideal gas | A5.21 |
| imaging detectors | F2.12 |
| imaging facility | F2.11 |
| incoherent neutron scattering cross section | A4.14 |
| incoherent scattering | A5.15, D7.2, E7.13 |
| index of refraction | D2.4, D2.10 |
| inelastic mean free path | F3.8 |
| inelastic scattering | A5.11, E7.8, E7.9, E7.10, E7.11, E7.12, E7.13, E7.14, E7.15 |

| | |
|--|--|
| information technology | E4.2 |
| inherent structure | E9.20 |
| integral dynamic measurement | E8.12 |
| inter diffusion | B3.3 |
| interface | E5.2, E5.9 |
| interface | D2.2, D2.5, D2.9, D2.11 |
| interference scattering | C7.37 |
| intermediate scattering function | A5.16, D8.3, D8.5, D8.10 |
| intermediate state | D4.15 |
| internal dynamics | E8.14 |
| Ir20Mn80 | E5.9 |
| iron | D9.5, D9.6, D9.7, D9.8, D9.11, D9.12, D9.13, D9.14, D9.17, E7.9, E7.10, E7.15, E7.16 |
| iron based superconductors | A1.35 |
| Ising model | D6.9-D6.12 |
| isomorphous replacement | E10.14 |
| isotope exchange | D1.18 |
| IXS | D4.8 |
| - J - | |
| Jahn-Teller distortion | D5.12 |
| Jahn-Teller effect | B5.17 |
| Jülich Centre for Neutron Science JCNS | A1.38 |
| jump diffusion | D7.12, E7.15, E8.11 |
| jumps between two sites | D7.10 |
| - K - | |
| Kauzmann | E9.1, E9.2, E9.3, E9.10, E9.11, E9.15, E9.18 |
| kinematical scattering | A3.2 |
| Klein-Nishina formula | A4.5 |
| Kramers-Kronig relation | A4.10, D11.4 |
| - L - | |
| L ₃ -phase | B2.3, B2.4 |
| LaAlO ₃ | E5.14 |
| Lamb-Mössbauer factor | D7.4 |
| lamellar phase | B2.4 |
| lamor precession | D8.3 |
| Lande factor | B5.15 |
| Langevin equation | B6.7 |
| LaNiO ₃ | E5.14 |
| large scale domain motions | E8.13 |
| Larmor diffraction | C6.8 |
| Larmor precession | C6.2 |
| lateral correlations | D2.30 |
| lattice vibrations | D4.2 |
| Laue | A1.2 |
| Laue function | A1.12 |
| Laue point | A3.7 |
| layer | D2.2, D2.15, D2.34 |
| LCLS (Linac Coherent Light Source) | C8.5 |
| length scale, mesoscopic | B2.2 |
| Lennard-Jones | E9.12, E9.14, E9.15 |
| lens | D1.7 |

| | |
|---|---|
| lensless imaging | D10.7 |
| level surfaces | B2.15, B2.16, B 2.32 |
| libration | D7.12 |
| ligand field | B5.3 |
| light scattering | D7.2 |
| Linac Coherent Light Source, LCLS | D10.2 |
| linear polarization | D11.10 |
| liquid | A5.9 |
| local disorder | D5.12 |
| local fluctuations | E7.10, E8.10 |
| local structure | D5.2, D5.3, D5.4, D5.8, E7.10 |
| localized motion | D7.6 |
| lone pairs | E4.11 |
| Lorentz correction | D3.11 |
| Lorentzian | D7.6 |
| low-spin complex | B5.10 |
| LuFe2O4 | E4.18 |
| - M - | |
| macromolecules | B2.2, B6.2 |
| magnetic dichroism | F3.29 |
| magnetic excitations | D4.2 |
| magnetic field (longitudinal) | D8.5 |
| magnetic fluids | E6.2, E6.3, E6.4, E6.10, E6.11, E6.12, E6.15, E6.16 |
| magnetic form factor for neutron scattering | A4.16 |
| magnetic interaction potential | C6.14 |
| magnetic linear dichroism | F4.26 |
| magnetic neutron scattering | A4.15, D6.15, D6.16, D6.20 |
| magnetic order | D11.8 |
| magnetic scattering | A1.19, E6.8, E6.9, E6.14, E6.15 |
| magnetic scattering length density | D2.22, D2.27 |
| magnetic structure | C7.2, C7.11, C7.19, C7.20, C7.21, C7.22, C7.23, C7.24, C7.25, C7.26, C7.27, C7.29, C7.30, C7.31, C7.32, C7.33, C7.36, C7.37 |
| magnetic x-ray scattering | A4.11, C7.2, C7.4, C7.13, C7.14, C7.19, C7.29, C7.37 |
| magnetism | D3.29, D9.13 |
| magnetite | E4.18 |
| magnetoelectric coupling | E4.9 |
| magnon | B4.14 |
| Markovian process | B6.6 |
| master equation | D7.6 |
| maximum entropy principle | B6.4 |
| mean relaxation time | D7.6 |
| mean squared displacement | B3.2, B3.6, B3.8, B6.7, B6.12, D3.6, D7.4, E3.8, E3.13, E9.3, E9.4 |

| | |
|--|--|
| mean-field theory | D6.4 |
| membrane bending rigidity | B2.9 |
| membrane fluctuations | B2.9, B2.12 |
| membrane model | B2.9 |
| membrane saddle-splay modulus | B2.9 |
| mesoscopic length scales | D1.2 |
| MF-RAM | E4.10 |
| micellar phase | B2.3 |
| micelles | B2.3, E5.3 |
| microbunching | C8.5 |
| microemulsion, near-wall structure | B2.29 |
| microemulsions | B2.6 |
| microemulsions (scattering intensity) | B2.14, B2.17, B2.22, B2.24, B2.28 |
| microemulsions (bicontinuous, lamellar) | D1.2, D8.9 |
| microscopy | D10.2 |
| Miller index | A5.8 |
| mixed crystal | D3.21 |
| mode coupling theory | E9.3, E9.12, E9.14, E9.15, E9.18, E9.20 |
| molecular dynamics | E7.15, E9.12 |
| molecular replacement | E10.15 |
| moment direction | C7.19, C7.22, C7.24, C7.26, C7.27, C7.33 |
| momentum distribution curve | F3.18 |
| momentum transfer | A5.3 |
| monatomic linear chain | B4.5 |
| monochromator | C3.4 |
| Monte Carlo simulations | B2.19 |
| mosaic crystal | C3.4 |
| Mössbauer spectroscopy | D9.2, D9.3, D9.4, D9.5, D9.11 |
| Mott-Bethe formula | A4.12 |
| multiferroic | E4.2 |
| multiple scattering | A3.2, C7.21, C7.22, C7.23, D3.10 |
| multiplexing | D4.7 |
| myoglobin | E10.17 |
| - N - | |
| nano-crystalline | D9.11 |
| nanoparticle structure | D5.11, D5.12 |
| neutron diffraction | D3.7 |
| neutron guide | C1.13, C3.3 |
| neutron magnetic moment | C6.2 |
| neutron polarisation | C3.6, C6.4 |
| neutron protein crystallography | E10.10 |
| neutron scattering | A4.13 |
| neutron spectrum | C1.2, C1.7, C1.12 |
| Neutron Spin Echo Spectroscopy (NSE) | C3.25, C6.8, D8.4, E3.9 |
| NEXAFS (Near-Edge X-ray Absorption Fine Structure) | F4.4 |
| nonresonant magnetic | C7.2, C7.7, C7.10, |

| | |
|---|--|
| scattering | C7.19, C7.27, C7.29, C7.34 |
| non-volatile memory | E4.3 |
| normal coordinates | B4.4 |
| normal mode analysis | E3.23 |
| normal modes | E8.7 |
| nuclear forward scattering (NFS) | D9.2, D9.3, D9.4, D9.5, D9.6, D9.8, D9.9, D9.10, D9.16, D9.17 |
| nuclear inelastic scattering (NIS) | D9.2, D9.3, D9.6, D9.8, D9.9, D9.15 |
| nuclear neutron scattering | D6.16 |
| nuclear quasi-elastic scattering (NQES) | D9.3., D9.15 |
| nuclear reactions | C1.3, C1.5, C1.6, C1.7 |
| nuclear reactor | C1.3, C1.5, C1.8, C1.10, C1.11 |
| nuclear resonance scattering (NRS) | D9.2, D9.4, D9.5, D9.14, D9.15, D9.16, D9.17, E7.8, E7.9, E7.11, E7.12 |
| nuclear scattering | A1.19 |
| numerical simulation | E9.1, E9.12, E9.13, E9.14, E9.18 |
| - O - | |
| object support | D10.9 |
| off-specular scattering | D2.4, D2.5, D2.7, D2.34 |
| oil-water-surfactant mixtures | B2.6 |
| One-Step Model | F3.11 |
| optic | B4.6 |
| optical interband transitions | F3.40 |
| optical spin orientation | F3.26 |
| optical theorem | D11.4 |
| orbital magnetic moment | F4.24 |
| orbital moment quenching | B5.11 |
| orbital order | D11.12 |
| orbital ordering | E5.14 |
| order parameter | D6.4, D6.8, D6.9, D6.19, D6.21 |
| Oseen tensor | B6.9 |
| osmotic pressure, sponge phase | B2.21, B2.23 |
| oversampling | C8.7 |
| - P - | |
| pair correlation function | A1.11, A1.26, A5.5 |
| partial PDF | D5.8 |
| partition function | B6.5 |
| PC-RAM | E4.22 |
| pendulum solutions | A3.15 |
| perovskite | D3.23 |
| phase change materials | E4.2 |
| phase contrast imaging | F2.17 |
| phase plate | C7.12, C7.16, C7.17, C7.18, C7.29, C7.27 |
| phase problem | A1.9 |
| phase retrieval | D10.2 |
| phase transition | D3.16, D6.2-D6.5, D6.7-D6.11, D6.13, |

| | |
|------------------------------------|--|
| | D6.17-D6.19, D6.21-D6.24 |
| phonon | B4.4, E7.15, E7.16 |
| phonon density of states | B4.8, E7.8 |
| phonon dispersion curves | B4.5, E7.11 |
| photoelectron diffraction | F3.23 |
| photoelectron spectroscopy | F3.2 |
| photosystem I | D10.17 |
| plateau modulus | E3.10 |
| pnictide | D3.29 |
| poincaré-Stokes vector | C7.11, D11.11 |
| polar nanoregions | D5.12 |
| polarization analysis | C7.11, C7.16, C7.20, C7.24, C7.25, C7.27, D11.10 |
| polarization analysis uniaxial | C6.9, C6.16 |
| polarization correction | D3.10 |
| polarization scan | D11.10 |
| polarized neutron scattering (PNR) | E5.12 |
| polarized neutrons | D2.22, D2.26, D8.5; E6.10, E6.14 |
| polyethyleneoxide (PEO) | B2.25 |
| polyethylenepropylene (PEP) | B2.25 |
| polymer | B6.2, D1.2, D1.14, E7.13, E7.14 |
| polymer diffusion coefficient | E3.6 |
| polymer dynamics | B6.2 |
| polymer melt dynamics | C3.29 |
| polymer solutions | B2.2 |
| polymer viscosity | E3.5 |
| polymer-induced bending rigidity | B2.26 |
| polymerized membranes | B2.20 |
| polymers, mushroom regime | B2.26 |
| polymers, overlap concentration | B2.26 |
| Porod scattering | D1.13 |
| powder average | D7.11 |
| powder diffraction | C3.8, D3.4, D3.14 |
| preaveraging approximation | B6.10 |
| precession | C6.20 |
| preferred curvature | B2.9 |
| Prigogine-Defay ratio | E9.1, E9.11, E9.12, E9.13, E9.14, E9.18, E9.19 |
| primary structure | E10.4 |
| protein | E10.2 |
| protein crystal | E10.9 |
| protein crystallography | E10.8 |
| protein data bank | E8.2 |
| protein dynamics | E8.8 |
| protein folding | E10.8 |
| protein internal dynamics | E3.23 |
| protein rigid body rotation | E3.23 |

| | |
|--|--|
| p-spin model | E9.14 |
| pump-and-probe experiments | C8.8 |
| pump-probe | E8.10 |
| PI-Flippers | D8.5, D8.6 |
| - Q - | |
| quantum beats | D9.4, D9.16 |
| quantum phase transitions | D6.18-D6.21 |
| quarternary structure | E10.8 |
| quasicrystal | B1.13 |
| quasielastic neutron scattering (QENS) | D7.2 |
| quasi-harmonic | B4.10 |
| - R - | |
| radiation damage | D8.13 |
| Ramachandran plot | E10.7 |
| random phase approximation | D1.17 |
| Rashba interaction | F3.27 |
| Rayleigh scattering | A4.4 |
| real-time imaging | F2.15 |
| reciprocal lattice | A5.9 |
| reduced dimensions/dimensionality | E5.2 |
| reflection | D2.3 |
| reflection coefficient | A3.17 |
| reflectivity | D2.9, D2.15, D2.20, D2.25 |
| refraction | D2.3 |
| renormalization | B2.12 |
| renormalized bending rigidity | B2.12 |
| replica | E9.3, E9.15, E9.18 |
| resistive oxide | F3.34 |
| resolution | A1.9, A1.30, D7.7, E10.12 |
| resolution (spatial, temporal) | D10.2 |
| resonance enhancement | D11.7 |
| resonance exchange scattering | D11.7 |
| resonant bonding | E4.5 |
| resonant magnetic scattering | C7.2, C7.4, C7.5, C7.6, C7.10, C7.11, C7.21, C7.23, C7.25, C7.28 |
| resonant scattering | D4.15, D11.3 |
| resonant scattering amplitude | D11.5 |
| Reverse Monte-Carlo | D5.9 |
| rigid body refinement | E8.5 |
| rocking curve | A3.18 |
| rotation of molecules | D7.11 |
| rotational constant | D7.11 |
| rotational diffusion | E8.14 |
| roughness | D2.5, D2.15, D2.31 |
| Rouse-model | B6.6, D8.8, E3.7 |
| - S - | |
| saddle splay modulus | B2.9, D8.9 |
| saddle-shaped membrane geometry | B2.4, D2.16 |

| | |
|--|--|
| salt bridge | E10.5 |
| SANS | C3.12, E8.4 |
| SASE (Self-Amplified Spontaneous Emission) | C8.3 |
| saturation length | C8.7 |
| sax | E8.4 |
| scaling | D6.10-D6.15, D6.18-D6.20 |
| scattering factor | A3.8 |
| scattering function | D4.12, D6.15-D6.17, D6.19 |
| scattering intensity, microemulsions | B2.14, B2.17, B2.22, B2.24, B2.28 |
| scattering law | A1.11, A1.26 |
| scattering length | A5.3, D3.7 |
| scattering length density | A5.5 |
| scattering length density | D2.10, D2.15, D2.18, D2.27 |
| scattering theory | A4.1 |
| scattering vector | A1.5, A5.3 |
| scattering wavevector | A4.8 |
| secondary relaxation | E9.2, E9.9, E9.15, E9.16, E9.17, E9.18 |
| secondary structure | E10.5 |
| seeding | C8.13 |
| self correlation function | D7.2, E7.13 |
| self diffusion (short-time, long-time) | B3.2, B3.3, D8.14, D8.18 |
| self-avoiding surfaces | B2.20 |
| self-organization | E5.3 |
| self-seeding | C8.13 |
| semiflexible polymer | B6.6 |
| SESANS | C3.13 |
| Shannon, Shannon sampling theorem | D10.5 |
| shrinkwrap | D10.12 |
| Shull | A1.3 |
| Siegbahn | F3.3 |
| single chain dynamic structure factor | E3.14 |
| single crystal | C3.8 |
| single crystal diffraction | D3.4, D3.19 |
| single-particle diffraction | D10.20 |
| small angle neutron scattering under grazing incidence | B2.29 |
| small angle scattering | A1.31 |
| small-angle neutron scattering | E6.2, E6.7, E6.15 |
| soft matter | B2.2 |
| soft mode | B4.9, D6.21-D6.23 |
| space group | B1.10 |
| spallation | A1.16 |
| spallation source | C1.6, C1.8, C1.9, C1.10, C1.11 |
| spatial resolution | F2.8 |
| speckle pattern | D8.14 |
| spectral density function | F3.13 |
| specular reflectivity | D2.3, D2.5, D2.9, D2.15, D2.26, D2.34, D2.36 |

| | |
|---|---|
| spherical | C6.9, C6.16 |
| spherical model | D6.10-D6.12 |
| spin and orbital angular momentum | C7.27, C7.29, C7.37 |
| spin anisotropy | B5.14 |
| spin magnetic moment | F4.24 |
| spin-filter barrier | F3.35 |
| spinodal decomposition | B3.24 |
| spin-orbit coupling | F3.4, F4.21 |
| spin-spiral ferroelectricity | E4.15 |
| spintronics | E5.2 |
| sponge phase | B2.3, B2.4, B2.22 |
| sponge phase, osmotic pressure | B2.21, B2.23 |
| spontaneous curvature | B2.9 |
| stacking faults | D5.9 |
| static scattering | A5.2 |
| statistical physics | B6.2 |
| steric membrane interaction | B2.4 |
| stochastic force | B6.6 |
| Stokes equation | B6.20 |
| stretched exponential function | D7.7 |
| stretching | E9.1, E9.6, E9.7, E9.8, E9.9 |
| structural dynamics | D9.2, D9.5, D9.8 |
| structure factor | A5.4, D3.6 |
| structure refinement | D3.12 |
| sum rules | F4.23 |
| superconductivity | D3.23 |
| superconductor | B1.14 |
| supercrystal | D2.31 |
| superlattice reflection | D11.15 |
| supermirror | C1.13 |
| superparamagnetism | E6.5 |
| surface | D2.2, D2.3, D2.34 |
| surface tension | B2.9 |
| surfactants | B2.2 |
| swedge method | F3.36 |
| symmetry | B1.7 |
| synchrotron radiation | A1.14, C7.2, C7.9, C7.10, C7.12, C7.13, C7.14, C7.15, C7.30 |
| synchrotron radiation based perturbed angular correlation (SRPAC) | D9.2, D9.3, D9.6, D9.16, D9.17, F2.3 |
| - T - | |
| tagged-particle dynamics | D7.2 |
| TbMnO ₃ | E4.13 |
| temperature (high/low) | D9.2, D9.6, D9.7, D9.8, D9.9, D9.10, D9.12, D9.13, D9.14, D9.15, D9.16, D9.17 |
| Templeton | D11.4 |
| ternary amphiphilic systems | B2.6 |
| tertiary structure | E10.7 |
| Teubner-Strey scattering | B2.17 |
| thermal neutrons | C1.7, C1.8, C1.9 |
| thin films | D9.14, E5.1 |

| | |
|--|--|
| Thomson scattering | A1.18, A4.5 |
| three axis spectrometer | C3.17, D4.6, E7.11 |
| Three-Step Model of Photoemission | F3.6 |
| tilt order | D11.12 |
| time average | D8.15, D8.11 |
| time-delay holography | D10.14 |
| time-of-flight spectrometer | C3.18, D4.9, D7.2 |
| tomography | F2.14 |
| topographic mapping | D10.21 |
| topology | B2.17 |
| topology fluctuations | B2.10 |
| total reflection | C6.4, D2.13, D2.24, D2.36 |
| total scattering | D5.2, D5.3, D5.4, D5.5, D5.6, D5.7, D5.9, D5.10, E7.11 |
| total scattering cross section | A4.2 |
| total scattering structure function | D5.2 |
| TPMS | B2.11, B2.29 |
| transition matrix element | F3.5 |
| translational diffusion | E8.14 |
| transmission | D2.3, D2.11 |
| transmission coefficient | A3.14 |
| triangulated surfaces, Monte Carlo simulations | B2.19 |
| triply periodic minimal surfaces | B2.11, B2.29 |
| tube mode | E3.11 |
| twinning | D3.24 |
| twist grain boundary | B2.29 |
| two-beam approximation | A3.6 |
| - U - | |
| undulator | C8.2 |
| universal memory | E4.2 |
| universality | D6.8-D6.10, D6.13 |
| unoccupied density of states | F4.20 |
| uphill diffusion | B3.29 |
| USANS | C3.12 |
| - V - | |
| van Cittert-Zernike theorem | D10.13 |
| van Hove correlation function | A5.17 |
| variational approach | B2.17 |
| velocity of sound | B4.6, E7.10 |

| | |
|--|-------------------------------|
| velocity selector | C3.5 |
| vesicle | B2.10 |
| vibrational density of states | D7.4 |
| vibronic interactions | B5.17 |
| Vogel-Fulcher | E9.1, E9.2, E9.3, E9.7, E9.10 |
| - W - | |
| water-surfactant mixtures | B2.3 |
| white noise | B6.6 |
| whole-particle structure modeling | D5.11 |
| wide angle diffraction | A1.31 |
| Wiener-Khinchine theorem | A5.30 |
| WLF | E9.7 |
| work function | F3.2 |
| - X - | |
| XANES (X-ray Absorption Near Edge Structure) | F4.4 |
| XPD | F3.23 |
| X-ray absorption spectroscopy (XAS) | E5.16 |
| x-ray crystallography | E10.9 |
| x-ray crystallography, nanocrystallography | D10.2 |
| x-ray diffraction | D3.7, D10.7 |
| x-ray edges | A4.9 |
| X-ray fluorescence | F4.7 |
| x-ray photon correlation spectroscopy (XPCS) | D8.11 |
| x-ray pulse | D10.14 |
| x-ray scattering | A4.8 |
| x-ray source | F2.3 |
| x-ray spectrum | F2.9 |
| x-ray standing wave | F3.36 |
| XSW | F3.36 |
| X-Y model | D6.9-D6.13 |
| xyz | C6.9, C6.7 |
| - Y - | |
| Yoneda effect | D2.13, D2.34, D2.36 |
| - Z - | |
| Zilman-Granek model | D8.10 |
| Zimm model | B6.9 |
| - 2-D detector (Pilatus) | |
| 2-D detector (Pilatus) | D8.12, D8.16 |
| 8-N rule | E4.6 |